

# Ethnobotanical and phytopharmacological notes on *Cymbopogon citratus* (DC.) Stapf

## Bibliographic Review

M. MACHRAOUI<sup>1\*</sup>, Z. KTHIRI<sup>1</sup>, M. BEN JABEUR<sup>1</sup>, W. HAMADA<sup>1</sup>

<sup>1</sup> Plant Production Department, Laboratory of Genetics and Plant Breeding, National Institute of Agronomy (INAT), 43, Av Charles Nicolle, 1082 Tunis, Tunisia

\*Corresponding author: machraoui.myriam@gmail.com

**Abstract** – *Cymbopogon citratus* (DC) Stapf is an herbal plant belonging to Poaceae. Commonly known as lemongrass, this plant is used in traditional and modern medicine to cure different diseases, as it has anti-inflammatory and anti-oxidant properties. Studies have shown that *Cymbopogon citratus* (DC) has interesting uses in agronomy as a natural pesticide and insecticide. Its antifungal and antimicrobial activities make it a plant of interest in many fields. Essential Oils, aqueous extracts, phenolic compounds and other extracts from this plant are industrially and economically important. They are used in perfumery, cosmetics and pharmaceuticals. The chemical composition of the *Cymbopogon citratus* (DC) essential Oil showed different molecules responsible of its therapeutic properties such as citral,  $\beta$ -myrcene, limonene, linalool, geranyl acetate, saponins, neryl acetate and tannins. These compounds have considerable antifungal, antibacterial, allopathic, and anti-carcinogenic activities. The following review is a summary of the ethnopharmacological use of *C. citratus*, but also its chemical composition and different biological activities.

**Keywords:** *Cymbopogon citratus*, citral, essential oil, antifungal activity, antibacterial activity.

### 1. General presentation of *Cymbopogon citratus* (D.C):

*Cymbopogon citratus* (DC.) commonly known as Lemongrass, Indian Lemongrass or Lemongrass from Madagascar is an aromatic herb belonging to the Poaceae family (Figure 1) This family is indeed widely distributed counting more than 635 genus and 9000 species. The *Cymbopogon* genus has for itself more than 140 cultivated species, 52 of them are located in Africa, 45 in India, 6 in Australia, 6 in South America, 4 in Europe, 2 in North America and the rest in South Asia (Suman et al., 2004). The rest of the species is distributed between Central America, South America, Africa and other tropical regions (Gagan et al., 2011).

Originated from southern India and Sri Lanka, *Cymbopogon citratus* (DC.) grows nowadays spontaneously all over the world, especially in the tropical subtropical and Savannah regions (Negrelle and Gomes, 2007). In West Africa, it has long been cultivated for its medicinal properties. In Tunisia, *Cymbopogon citratus* (DC.) is an introduced species cultivated for decoration, therapeutics and insecticides uses. Indeed, it is grown in gardens especially for its repelling effect of insects like mosquitoes. The essential oil extracted from the lemongrass is used in local industrial products against cockroaches, flies and mosquitoes (Dhaou et al., 2010).

### 2. Taxonomy classification


*Cymbopogon citratus* (DC.) is a herb belonging to the Kingdom of Plantae, the Phylum of Spermatophyta (seed plants), the super-branching of Magnoliophyta (flowering plants), the class of Liliopsida (Monocotyledons), the order of Poales, the family of Poaceae (herbaceous), the genus is *Cymbopogon* and the species is *citratus* (Karunamoorthi et al., 2010).

### 3. Morphological and botanical description

*Cymbopogon citratus* (DC.) Stapf (Figure 1) is a perennial aromatic herb that grows as dense clumps with no ramification (Fig1.A). The total length of this plant can reach 2 meters while its width goes up to 1.2 m. It has short rhizomes which also are its way of multiplication (Fig1.B). The leaves are green, erect, flat, linear in shape and closed at the base (Figure 1.C). The majority emerges directly from the

ground without stem. Their length exceeds 1m while the width varies from 5 to 15 mm. The leaves give off a characteristic lemon flavor, once pressed by hand or crushed. The upper side is glabrous and whitish with ligaments 4 to 5 mm long.

The glumes are equal to subequal. The lower glume is lance-shaped with an acute apex, while the upper glume is lanceolate 4.3 to 4.5 cm long with a rib. *C. citratus* rarely gives flowers. The species identified until now do not show flowers. The inflorescence: *C. citratus* has erect inflorescences with a length of 30 to 60 cm. *C. citratus* is generally propagated by seedling or by tuft division (Negrelle and Gomes, 2007; Gagan et al., 2011).



**Figure 1.** *Cymbopogon citratus*: A: the aerial part, B: The root system, C: one leaf

#### 4. Common names

*Cymbopogon citratus* (DC.) has different names depending on the country in which it grows. In France and Tunisia, it's called "Citronnelle" or "Herbe Citron". In USA and England, English, they call it "Lemongrass" or "Lemon Grass" which is also the common name in India and Egypt. In Brazil it's called "Capim-cidrao" or "Capim-santo", in Ethiopia "Tej-sar". For the rest of the countries, the most used names are the following: Sereh (Indonesia), Cimbopogone (Italy), Sakumau (Malaysia), Zacate limon (Mexico City), Citrongrass (Sweden), Ta-khrai (Thailand), Limon out (Turkey) (Gagan et al., 2011).

#### 5. Ethnopharmacological uses of *Cymbopogon citratus* (DC.)

*C. citratus* has always been grown all over the world for decoration but also for medical purposes (Tarkang et al., 2012). In traditional medicine, it was used for its virtues on human health depending on the country or even the continent where it is located. In Egypt, dried leaves were boiled and orally administered as a renal anti-spasmodic and diuretic. Leaves and essential oil (EO) showed significant results in treating cough, fever, vomiting, headache and even insomnia and depression (Oloyede, 2009). In Indonesia and Malaysia the whole plant is boiled and the liquid obtained is administered orally for its emmenagogue effect (stimulation of the pelvic blood flow and uterine region). In the United States of America, the boiled liquid is used as an external lotion to heal wounds and bone fractures, especially in the Minnesota region (Spring, 1989). In South America, precisely in Brazil, tea obtained from the leaves of *C. citratus* is drunk for its antispasmodic, anti-inflammatory and analgesic effects (Fransisco et al., 2013). In India: the plant itself of *C. citratus* is grown in gardens to ward off snakes. Indian also give children lemongrass tea with slices of lemon to heal stomatitis. A few drops of the EO diluted in warm water is administered orally to treat gastrointestinal problems. In Africa, it is used as an antitussive, antiseptic, sudorific and to treat back pain.

Lemongrass leaves are also brought to a boil with the leaves of bamboo (*Bambusa vulgaris*) and ginger to reduce fever. In Nigeria and Ghana, decoctions of *C. citratus* leaves with *Cassia occidentalis* and/or key lime (*Citrus aurantifolia*) is used to treat malaria especially for pregnant women. Patients take one cup per day until complete healing. Chinese treat colds and oral bleeding with *C. citratus* essential oil (Negrelle et al., 2007; Tchoumboungnan and al., 2005).

Studies have also shown that lemongrass has allelopathic properties, which means it can be used as a herbicide (Dudai et al., 1999). Lemongrass essential oil concentrated at 10% in an aqueous suspension has a total herbicidal effect on many weed species such as *Digitaria horizontalis*, *Sorghum halepense*,

*Bidens pilosa*, *Euphorbia heterophylla* and *Raphanus raphanistrum* (Valarini et al., 1996). Even animals are treated with *C. citratus*. In Brasil, the tea made out of leaves and roots is given as a remedy for colic. The oil showed its efficiency against body odors (Lachman-White and al., 1992). Macerate leaves reduces swelling, mosquito's bites, wounds and eczema (Asase et al., 2010). Moreover, *C. citratus* is known for its domestic use. In aerosols, deodorants, *C. citratus* is used as insect repellent. Roots extracts are added to perfumes, shampoo, soaps and cosmetics. It is also used as a household detergent (Ritter et al., 2012).

## 6. Pharmacological uses of *C. citratus*

Different pharmaceutical and medical studies showed that *C. citratus* essential oil has antimicrobial, antiparasitic and antifungal activities (Tzortzakis and Economakis, 2007). Inhibiting the proliferation of cancer cells, lemongrass EO is also an antioxidant, anti-tumoral and anti-inflammatory. This EO helps reducing blood pressure and protect the cardiovascular and gastric system (Manosroi et al., 2006).

## 7. Chemical composition of *C. citratus* essential oil

The essential oil of *C. citratus*, is obtained by hydrodistillation. The yield varies according to the part of the plant from which the extraction took place. Generally speaking, the leaves yield varies from 0.28 to 1.4% (Kasali and al., 2001), given that the maximum value obtained is 3% from the dry and non-green leaves (Chisowa and al., 1998). The chemical compounds obtained from the roots are different from those obtained from the leaves. These molecules are classified in four groups

oxygenated monoterpenes, oxygenated sesquiterpenes, monoterpene hydrocarbons and hydrocarbons sesquiterpenes. Other components have also been identified from leaves and roots, including alkaloids, saponins, flavonoids, caffeic acids and sugars (Akhila, 2010). *C. citratus* is most of the times grown for its high content of Citral, a component that is used in cosmetic and chemical industries (Miean and Mohamed, 2001). Citral is also the active ingredient responsible for the antifungal and antimicrobial activity which accounts for 70% to 80% of the total yield obtained.

Previous studies of *C. citratus* have shown that the chemical composition also depends on the geographical origin. Moreover, the essential oil contains terpene carbons, alcohols, ketones, esters and aldehydes. Organic obtained with methanol and ethanol and aqueous extracts obtained with infusion of the leaves showed that *C. citratus* contains tannins, phenolic acids and flavonoid glycosides. *C. citratus* also contains vitamins A, C, E but also niacin, pyridoxine, riboflavin and folate. Concerning minerals, *C. citratus* contains sodium, potassium, manganese, zinc, calcium, magnesium, copper selenium, phosphorus and iron in addition to electrolytes, macronutrients (carbohydrates, proteins), and a small quantity of fat (Cheel and al., 2005). Other compounds were reported such as orientin, swertiajaponin and isoscoparin as well as other phytochemicals (Bahrati et al., 2013). Table 1 summarized the components found in *C. citratus* EO.

## 8. Biological activities of *Cymbopogon citratus* EO

### 8.1. Antimicrobial activity

The essential oil extracted from lemongrass leaves is widely used for its antimicrobial activity. Concerning the bacteria, *C. citratus* seems to be more efficient against Gram negative ones than Gram positive (Mayaud et al., 2008). Against *E. coli*, the EO showed over 80% of efficiency. It was even more active against *Enterococcus faecalis* and *Salmonella typhimurium* than two standard antibiotic which are erythromycin tetracycline (De Oliveira et al., 2013). *C. citratus* EO was more efficient than *Mentha arvensis* var. *piperacens* EO against enteropathogenic germs which can contaminate and affect food quality. Moreover, this EO inhibits the activity of *Salmonella enteritidis* responsible for deterioration of preserved meat. Patients who have been given 05ml/day of *C. citratus* was healed from *Helicobacter pylori*, the pathogen responsible of gastric ulcer. Its action in acid medium was bactericidal (Ohno et al., 2003). Generally speaking, the antimicrobial activities of *C. citratus* EO are mainly attributed to geraniol ( $\alpha$ -citral), neral ( $\beta$ -citral), 1, 8- cineole, *p*-cymene,  $\alpha$ - and  $\beta$ -pinene, limonene,  $\alpha$ -terpineol and camphene (Cimanga et al., 2002). The essential oil also exhibits good antibacterial activity, in particular against *Escherichia coli*, which is harmful to humans (Ogunlana et al., 1986).

**Table 1.** Natural compounds reported in *C.citratus* EO with possible amounts 16 18 190 192

Chemical composition of <i>C.citratus</i> Essential Oil: main components	References
<i>α</i> -citral (geranial) (29,4 - 60,3%)	(Ekpenyong et al.,2015)
limonen (3-5,4%)	
6-methyl-hepten-2-one (2,9 - 3,21%)	
citronella (0,2-1,32%)	
germacren-D (0,2 - 0,5%)	
, n-octanal (0,2%)	
1,8-cineol (0,2%)	
z-carveol (0,2%)	
$\gamma$ -cadinen (1,3%)	
<i>α</i> -terpinéol (0,29 - 9 %)	
$\beta$ -citronellal (0,2 - 0,7%)	(Nambiar et Matela, 2012)
$\beta$ -caryophylen (1,9%)	
<i>α</i> -Amorphene	
$\beta$ -Sesquiphellandrene	
Isolongifolene-4-5-9-10-dehydro Levo-	
<i>β</i> -elemene	
<i>α</i> -Bergamotene	
<i>t</i> -Cadinol	
<i>α</i> -citral (geranial) (29,4 - 60,3%)	(Bassolé et al., 2011)
limonen (3-5,4%)	
6-methyl-hepten-2-one (2,9 - 3,21%)	
citronella (0,2-1,32%)	
germacren-D (0,2 - 0,5%)	
, n-octanal (0,2%)	
1,8-cineol (0,2%)	
z-carveol (0,2%)	
$\gamma$ -cadinen (1,3%)	
<i>α</i> -terpinéol (0,29 - 9 %)	
$\beta$ -citral (neral) (21,39 - 40%)	(Asaolu et al.,2009)
Neryl acetate (2,0 - 6,3%)	
aromadendrene (0,6%)	
methyl heptone (1-1,4%)	
3-carene (0,1%)	
$\beta$ -myrcen (0,8 - 20%)	
tricyclen (0,2%)	
verbenone (0,2%)	
sabinol (0,5%)	
linalool (0,2 - 3,2%)	(Faruq, 1994)
<i>α</i> -cyclocitral (0,5%)	
genaryl acetate (0,2 - 3,6%)	
<i>α</i> -Muurolene	
d-Cadinene	
3-Undecyene	
3-carvomethenone	
$\gamma$ -Muurolene	
Dextro-carvone	
nerol acetate (10,8%)	(Tzortzakakis and Economakis, 2007)
geranial (3,25 -4%)	
<i>α</i> -pinen (0,4 - 1,1%)	
myrtanal (0,2%)	
terpinen-4-ol (0,5 - 3%)	
camphen (0,9 - 1,5%)	
eugenol (0,24 - 0,3%)	
borneol (0,2 - 3,7%)	(Akhila, 2010)
isoeugenol (0,5%)	
nerol acid (0,8%)	
caryophylen oxide (0,8%)	
$\beta$ -ocimen (0,3%)	
bomeol (1-2,16%)	
Germacrene-D	
( <i>E</i> , <i>E</i> )-Farnesal pimelyl	
Dihydrazide	
<i>α</i> -Gurjunene	

## 8.2. Antifungal activity

Pathogen resistance to a certain number of chemical treatments and target mutation is becoming a serious issue. For this particular reason, natural products are used today as a promising alternative. The essential oil extracted from lemongrass leaves is widely used for its antifungal activity. The EO obtained by hydrodistillation was effective against 42 microorganisms including 7 yeasts and 15 fungi (Negrelle and Gomez, 2007). These fungi include rice pathogens such as *Rhizoctonia solani* (Shimoni, 1993). Antifungal activities of the *C. citratus* oil can be attributed to the presence of a number of compounds among which citral,  $\beta$ -myrcene, linalool, and geraniol (Alvanioano et al., 2005). Studies showed that the mixture of geranium and neral, also called citral, that is responsible for the antifungal activity. Moreover, with the presence of myrcene with one of the two compounds, this activity is further enhanced (Di Pasqua et al., 2006).

*C. citratus* is also effective against fungi causing mold such as *Aspergillus niger*, *Aspergillus ochraceus*, *Alternaria alternata*, *Fusarium oxysporum* and *Penicillium roquefortii*, at a concentration going between 500 ppm and 700ppm (Helal et al., 2006). The EO also inhibits the production of aflatoxine B1, a toxin produced by *Aspergillus flavus* (Singh et al., 2010). The effective doses against these following yeasts were ranged between 0.062  $\mu$ l/ml and 10 $\mu$ l/ml: *C. albicans*, *Candida oleophila*, *Hansenula anomala*, *Schizosaccharomyces pombe*, *Saccharomyces cerevisiae*, *Saccharomyces uvarum* et *Metschnikowia fructicola*. Generally speaking, fungi were more sensitive to *C. citratus* EO than Yeasts (Irkin and Korukluoglu, 2009). This shows that this aromatic plant can be a perfect substitute to chemical treatments on plants, human and animals.

## 8.3. Insecticide activity

*C. citratus* has always been grown in gardens to repulse insects like cockroaches and mosquitoes. *In vitro* experiments on the common housefly, *Musca domestica* L. (Diptera: Muscidae) showed that Assay of oil against housefly larvae and pupae through contact toxicity assay showed lethal concentration (LC)<sub>50</sub> value of 0.41  $\mu$ l/cm<sup>2</sup> and of percentage inhibition rate (PIR) of 77.3 %, respectively (Kumar et al., 2013). While oil from *C. citratus* demonstrated complete larvicidal action on the yellow fever mosquito *Aedes aegypti* L. Twenty giving 100% mortality of at 200 ppm (Sosan et al., 2001). Against *Culex quinquefasciatus* and *Anopheles dirus*, 10% of *C. citratus* EO applied in soybean oil 1%, gave a 100% mortality rate 24 hours after the application. This shows the potentiality of *C. citratus* to be used as a natural insecticide instead of chemical products (Phasomkusolsil & Soonwera, 2011). Using also contact toxicity against *Tribolium castaneum* (Herbst.) (Coleoptera: Tenebrionidae), *C. citratus* EO was repellent at a dose of 0.021 ml/L after 4 hours of exposure. Its efficiency was much higher than the active ingredient IR3535 repellent found in a variety of insect repellent products used at the dose of 0.686 ml/L. This insecticide activity is mostly due to the presence of geraniol and neral in the chemical composition of the EO (Olivero-Verbel et al., 2010). The essential oil of lemongrass is also used against insects, especially mosquitoes, stopping certain phases of its development cycle. This has been proven for the mosquito *Aedes aegypti* (the tiger mosquito) vector of yellow fever (Osmani and Sighamony, 1980).

## 9. Medical uses and benefits on human health

The use of lemongrass in traditional medicine depends on the country where it is grown. However, tea or infusion prepared from dry leaves is used in broad spectrum treatment (Tarkang et al., 2012).

**Table 2.** Uses of *C. citratus* in traditional medicine

Country	General use of <i>C. citratus</i>	References
South Africa	Antitussive, antiseptic, sudorific, antirheumatic, treatment of back pain and sprains.	(Ekpenyong et al., 2015; Negrelle and Gomes EC, 2007)
Asia: India	Treatment of Gastrointestinal Problems	
Suriname	cough, asthma, bladder problems, wound management and headaches	
Indonesia	facilitate digestion, diuresis, sweating and used as an emmenagogue	
Brazil	colds, coughs, heart disease, urinary tract inflammations and allergies. It is also used to treat women neuropsychological problems	
Niger	antipyretic and antispasmodic.	

### 9.1. Regulation of the cardiovascular system

In traditional medicine, *C. citratus* is used for hypotension treatment. Tested on rats at different doses going from 1 to 20 mg/Kg, *C. citratus* EO induces hypotension, by reduction in vascular resistance caused by inhibition of the  $Ca^{2+}$  influx, and bradycardia probably due to an activation of cardiac muscarinic receptors (Moreira et al., 2010). Furthermore, aqueous extracts obtained from pulverized dry leaves of *C. citratus* were tested on cardiac rate and contractile force on the isolated hearts of rats. These extracts provoked significant Cardiac Rate reduction without altering the Contractile Force. This activity is due to the presence of alkaloids, tannins and flavonoid (Gazola et al., 2004). Another experiment was conducted on aqueous crude extracts of *C. citratus* showed its efficiency on the osmotic stability of human erythrocytes. The extracts protected the erythrocytes against hypotonic shock by a decrease of the value of the intensity of the lysis effect in NaCl solutions (De Freitas et al., 2008).

### 9.2. Anti-inflammatory properties

*C. citratus* is an aromatic herb that contains dietary components responsible for an anti-inflammatory activity. Infusion of dried leaves inhibits the lipopolysaccharide (LPS) induced by NO production and inducible NO synthase (iNOS) protein expression. These two later are involved in inflammatory cells activation and chronic inflammation. *C. citratus* its polyphenolic fractions, flavonoid, tannin and phenolic acid-rich fractions reduced iNOS protein levels and NO production especially in gastrointestinal diseases (Figueirinha et al., 2010). Moreover, polyphenols obtained from *C. citratus* extracts inhibited cytokine production on human macrophages. Chlorogenic acid (CGA), main phenolic acid of the infusion, is the agent responsible of this anti-inflammatory activity (Francisco et al., 2013). The EO was also shown effective against ileum inflammation in mice. It inhibits  $\beta 7$ -expression, responsible of lymphocytes migration (Watanabe et al., 2010).

### 9.3. Antioxidant activity

*C. citratus* has been used for medical purposes to treat pathogenesis neurological disorders. Containing antioxidant compounds, it helps reduce the oxidative stress important in the pathogenesis of several neurological diseases. Quercetin, gallic acid, quercetin and rutin are the phenolic compounds found in *C. citratus* extracts. These later protect against oxidative damage induced by various pro-oxidant agents that induce lipid peroxidation. Thus *C. citratus* could be considered an effective agent in the prevention of various neurological diseases associated with oxidative stress (Pereira et al., 2009). However, 1% of *C. citratus* dried powder added to pork raw meat, improves its conservation by significantly reducing Thiobarbituric Acid Reactive Substances (TBARS) (Olorunsanya et al., 2010). Methanolic, aqueous extracts as well as infusion and decoction of *C. citratus* showed significant activity in inhibiting the enzyme xanthine oxidase (XO) and lipid peroxidation in human erythrocytes. Isoorientin and orientin presented similar activities toward the DPPH (IC<sub>50</sub>: 9–10  $\mu$ M) reducing lipid peroxidation by 70% at 100  $\mu$ g/mL. Caffeic and chlorogenic acid were active superoxide anion scavengers with IC<sub>50</sub> values of 68.8 and 54.2  $\mu$ M, respectively, and a strong effect toward DPPH. Caffeic acid inhibited lipid peroxidation by 85% at 100  $\mu$ g/mL (Cheel et al., 2005).

### 9.4. Anti-carcinogenic and antimutagenic activities

Numerous studies have been carried out on the antitumor activity of citronella extracts. All of them indicated that *C. citratus* has a promising anticancer activity causing loss in tumor cell viability. Tested on Fischer rats, *C. citratus* extracts against hepatocarcinogenesis, ethanolic extracts at a dose going from 0.2% to 1.8% reduces the incidence of hyperplastic lesions and mammary murine epithelial cells apoptosis (Puatanachokchai et al., 2002). *In vitro*, polysaccharides extracted from *C. citratus* have shown anticancer activities due to the presence of (1-4) linked b-D-Xylofuranose moiety. These polysaccharides are also responsible of apoptosis in cancer cells *via* intrinsic pathway through the events of up-regulation of caspase 3, down-regulation of bcl-2 family genes followed by cytochrome *c* release (Thangam et al., 2014). In a cup of tea prepared with 1g of *C. citratus* leaves, 44.5 $\mu$ M of Citral induces apoptosis in hematopoietic cancer lines. The apoptosis was accompanied by DNA fragmentation and caspase-3-catalytic activity induction. However, citral apoptotic activity was dependent on the  $\alpha$ ,  $\beta$ -unsaturated aldehyde group. This anticancer effect was higher than a reference compound

called staurosporine (Dudai et al., 2005). Against solid and ascitic Ehrlich and Sarcoma-180 tumor models in mice, *C. citratus* showed a dose dependent effect going from 4.2 to 79 µg/ml. The highest cytotoxicity was noticed against 502713 (colon) and IMR-32 (neuroblastoma) cell lines with an IC<sub>50</sub> of 4.2 to 4.7 µg/ml. The total tumor inhibition with both ascitic and solid tumor forms of Ehrlich Ascites carcinoma needed doses of 97.34 and 57.83 respectively. Moreover, the EO tested on Sarcoma-180 solid tumor cells from animals showed condensation and fragmentation of nuclei typical of apoptosis (Sharma et al., 2009). Citral also showed activity against leukemic cells called P388 (Dubey and al., 1997). The ethanolic extracts of *C. citratus* also have anti-mutagenic activity (Vinitketkumnuen et al., 1994). Myrcene, showed an antimutagenic effect specifically on mammary cells. While α-limonene and geraniol have been able to inhibit liver and intestinal mucosal membrane in mices (Zheng and al., 1993). It has also been shown that juices extracted from *C. citratus* leaves, showed inhibiting activity containing inhibitors of cutaneous tumors development (Lima et al., 1993).

#### **9.5. Blood glucose regulation and hypolipidemic effects**

Aqueous extract of *C. citratus* has hypoglycemic effects. A daily dose going from 125-500 mg per kilogramme tested on Wistar rats' lowers fasting plasma glucose (FPG) but also total cholesterol and lipid parameters such as high-density lipoprotein (HDL). And because this used dose did not show any type of toxicity, *C. citratus* can be used as a treatment against type 2 diabetes (Adeneye et al., 2007). Moreover, a treatment with *C. citratus* extracts during 4 weeks on diabetic rats lowers blood glucose, TG, T-chol and LDL levels. This same treatment induced body weight reduction. The EO showed hypocholesterolemic effect mediated by the regulatory enzyme, HMG-CoA reductase in post-transcriptional down-regulation (Middleton and Hui, 1983). This way, the EO inhibits the hepatic 3-hydroxy-3-methyl-glutaryl-coenzyme A (HMG-CoA) reductase that plays an important role in cholesterol synthesis (Elson et al., 1989).

#### **10. Toxicity of *C. citratus* EO**

Applied at doses generally higher than 1500 mg/kg body weight, the oils caused significant functional damages to stomach and liver of Wistar rats for consecutive 14 days (Fandohan et al., 2008). The herbal tea made out of dried leaves of *C. citratus* caused slight elevations of direct bilirubin and amylase on humans but without clinical manifestation (Leite et al., 1986). At an elevated dose of 2000mg/Kg of *C. citratus* EO tested on Wistar mices, symptoms like torpor, nose and eyelid bleeding were observed. These symptoms were due to stomach mucosa atrophy and hepatocytes necrosis (Costa et al., 2011). Concerning the plants, the aqueous extracts of *C. citratus* on *Allium cepa* bulb (onion) at doses going from 1 to 20% induced cytotoxic and genotoxic effects (Akinboro and Bakare, 2007). Aqueous extract tested on *Lactuca sativa* at a concentration of 30mg/ml caused chromosomal abnormalities and root cell death (Sousa et al., 2010). The major components causing toxicity in *C. citratus* are citral and β-myrcen. Limonen showed low toxicity on humans with a dose of 100 mg/Kg causing nausea, vomiting and diarrhea (Vigushin et al., 1998). At doses going from 125 mg/kg up to 1200 mg/kg, citral and β-myrcen become toxic on embryos. Growth is delayed and skeletal abnormalities is observed with an increase in the fetal spleen weight (Nogueira et al., 1995).

#### **11. Conclusion**

Scientific research conducted on *Cymbopogon citratus* helped understanding the beneficial effects on human, animals and plants. Its pharmacological properties explained the different uses of this plant in medicine as a digestive and cardiovascular regulator but also against infectious diseases and psychiatric disorders. The chemical composition of the EO, phenolic compounds and aqueous extracts showed the major compounds which are responsible of the antifungal, antimicrobial, antioxidant and anticancer activities. The different information detailed in this review on the benefits and uses of *C. citratus* make it a good candidate for further scientific experiments in other fields such as agronomy and pharmaceuticals.

## 12. References

- Adeneye, A. A., & Agbaje, E. O. (2007).** Hypoglycemic and hypolipidemic effects of fresh leaf aqueous extract of *Cymbopogon citratus* Stapf. in rats. *Journal of ethnopharmacology*, 112(3), 440-444.
- Akhila A.** *Essential oil bearing plants: The genus Cymbopogon*. Edited by: Anand Akhila [M]. Broca Raton, FL: CRC Press Taylor & Francis Group, 2010.
- Akinboro A, Bakare AA (2007)** Cytotoxic and genotoxic effects of aqueous extracts of five medicinal plants on *Allium cepa* Linn. *J Ethnopharmacol* 112: 470–5
- Alviano WS, Mendonca RR, Alviano DS, *et al.* Antimicrobial activity of *Croton cajucara* Benth linalool-rich essential oil on artificial biofilms and planktonic microorganisms [J]. *Oral Microbiol Immunol*, 2005, 20 (2): 101-105.
- Asaolu MF, Oyeyemi OA, Olanilokun JO.** Chemical compositions, phytochemical constituents and *in vitro* biological activity of various extracts of *Cymbopogon citratus* [J]. *Pakistan J Nutr*, 2009, 8(12): 1920-1922.
- Asase A, Akwetey GA, Achel DG (2010)** Ethnopharmacological use of herbal remedies for the treatment of malaria in the Dangme West District of Ghana. *J Ethnopharmacol* 129: 367–76
- Bassolé, I. H. N., Lamien-Meda, A., Bayala, B. O. L. C., Obame, L. C., Ilboudo, A. J., Franz, C., ... & Dicko, M. H. (2011).** Chemical composition and antimicrobial activity of *Cymbopogon citratus* and *Cymbopogon giganteus* essential oils alone and in combination. *Phytomedicine*, 18(12), 1070-1074.
- Bharti SK, Kumar A, Prakash O, et al.** Essential oil of *Cymbopogon citratus* against diabetes: validation by *in vivo* experiments and computational studies [J]. *Scientif Rep*, 2013a, 2(3):1-9 688. doi:10.4172/scientificreports.688
- Cheel, J., Theoduloz, C., Rodríguez, J., & Schmeda-Hirschmann, G. (2005).** Free radical scavengers and antioxidants from Lemongrass (*Cymbopogon citratus* (DC.) Stapf.). *Journal of agricultural and food chemistry*, 53(7), 2511-2517.
- Chisowa, E. H., Hall, D. R., & Farman, D. I. (1998).** Volatile constituents of the essential oil of *Cymbopogon citratus* Stapf grown in Zambia. *Flavour and Fragrance Journal*, 13(1), 29-30.
- Cimanga K, Apers S, de Bruyne T, et al.** Chemical composition and antifungal activity of essential oils of some aromatic medicinal plants growing in the Democratic Republic of Congo [J]. *J Essential Oil Res*, 2002, 14(5): 382-387.
- Costa CA, Bidinotto LT, Takahira RK, et al. (2011)** Cholesterol reduction and lack of genotoxic or toxic effects in mice after repeated 21-day oral intake of lemongrass (*Cymbopogon citratus*) essential oil. *Food Chem Toxicol* 49(9): 2268–72
- De Freitas, M. V., Rita de Cássia, M. N., da Costa Huss, J. C., de Souza, T. M. T., Costa, J. O., Firmino, C. B., & Penha-Silva, N. (2008).** Influence of aqueous crude extracts of medicinal plants on the osmotic stability of human erythrocytes. *Toxicology in Vitro*, 22(1), 219-224.
- De Oliveira TL, Soares R De A, Piccoli RH (2013)** A Weibull model to describe antimicrobial kinetics of oregano and lemongrass essential oils against *Salmonella enteritidis* in ground beef during refrigerated storage. *Meat Science* 93: 645–51
- Dhaou, S. O., Jeddi, K., & Chaieb, M. (2010).** Les Poaceae en Tunisie: systématique et utilité thérapeutique. *Phytothérapie*, 8(2), 145-152.
- Di Pasqua R, Hoskins N, Betts G, et al.** Changes in membrane fatty acids composition of microbial cells induced by addition of thymol, carvacrol, limonene, cinnamaldehyde, and eugenol in the growing media [J]. *J Agric Food Chem*, 2006, 54(7): 2745-2749.
- Dubey, N. K., Takeya, K., & Itokawa, H. (1997).** Citral: A cytotoxic principle isolated from the essential oil of *Cymbopogon citratus* against P388 leukaemia cells. *Current science*, 73(1), 22-24.
- Dudai, N., Poljakoff-Mayber, A., Mayer, A. M., Putievsky, E., & Lerner, H. R. (1999).** Essential oils as allelochemicals and their potential use as bioherbicides. *Journal of Chemical Ecology*, 25(5), 1079-1089.
- Dudai, N., Weinstein, Y., Krup, M., Rabinski, T., & Ofir, R. (2005).** Citral is a new inducer of caspase-3 in tumor cell lines. *Planta medica*, 71(05), 484-488.
- Elson CE, Underbakke P, Hanson E, et al.** Impact of lemongrass oil, an essential oil on serum cholesterol [J]. *Lipids*, 1989, 24 (8): 7677-7679.

- Ekpenyong, C. E., Akpan, E., & Nyoh, A. (2015).** Ethnopharmacology, phytochemistry, and biological activities of *Cymbopogon citratus* (DC.) Stapf extracts. *Chinese journal of natural medicines*, 13(5), 321-337.
- Fandohan, P., Gnonlonfin, B., Laleye, A., Gbenou, J. D., Darboux, R., & Moudachirou, M. (2008).** Toxicity and gastric tolerance of essential oils from *Cymbopogon citratus*, *Ocimum gratissimum* and *Ocimum basilicum* in Wistar rats. *Food and chemical Toxicology*, 46(7), 2493-2497.
- Faruq MO,** TLC technique in the component characterizations and quality determination of Bangladeshi lemongrass oil (*Cymbopogon citratus*)(DC) Stapf.) [J]. *Bangladesh J Sci Ind Res*, 1994, 29(2): 27-38.
- Figueirinha, A., Cruz, M. T., Francisco, V., Lopes, M. C., & Batista, M. T. (2010).** Anti-inflammatory activity of *Cymbopogon citratus* leaf infusion in lipopolysaccharide-stimulated dendritic cells: contribution of the polyphenols. *Journal of Medicinal Food*, 13(3), 681-690.
- Francisco, V., Costa, G., Figueirinha, A., Marques, C., Pereira, P., Neves, B. M., ... & Batista, M. T. (2013).** Anti-inflammatory activity of *Cymbopogon citratus* leaves infusion via proteasome and nuclear factor- $\kappa$ B pathway inhibition: contribution of chlorogenic acid. *Journal of ethnopharmacology*, 148(1), 126-134.
- Gagan Shah, Richa Shri, Vivek Panchal, Narender Sharma, Bharpur Singh and A. S. Mann (2011)** Scientific basis for the therapeutic use of *Cymbopogon citratus*, stapf (Lemon grass). *J Adv Pharm Technol Res*. 2011 Jan-Mar; 2(1): 3–8.
- Gazola, R., Machado, D., Ruggiero, C., Singi, G., & Alexandre, M. M. (2004).** Lippia alba, Melissa officinalis and *Cymbopogon citratus*: effects of the aqueous extracts on the isolated hearts of rats. *Pharmacological research*, 50(5), 477-480. Miean KH, Mohamed S. Flavonoid (Myricitin, Quercetin,
- Helal GA, Sarhan MM, Abu Shahla ANK, et al. (2006)** Effects of *Cymbopogon citratus* L.essential oil on the growth, lipid content and morphogenesis of *Aspergillus niger* ML2-strain. *J Basic Microbiol* 46: 456–69
- Irkin R, Korukluoglu M (2009)** Effectiveness of *Cymbopogon citratus* L. Essential oil to inhibit the growth of some filamentous fungi and yeasts. *J Med Food* 12(1): 193–7.
- Kaempferol, Luteolin, and Apigenin)** content of edible tropical plants [J]. *J Agric Food Chem*, 2001, 49(6): 30106-30112.
- Karunamoorthi K, Ilango K, Murugan K (2010)** Laboratory evaluation of traditionally used plant-based insect repellent against the malaria vector *Anopheles arabiensis* Patton (Diptera: Culicidae). *Parasitol Res* 106:1217–23
- Kasali, A. A., Oyediji, A. O., & Ashilokun, A. O. (2001).** Volatile leaf oil constituents of *Cymbopogon citratus* (DC) Stapf. *Flavour and Fragrance Journal*, 16(5), 377-378.
- Kumar, P., Mishra, S., Malik, A., & Satya, S. (2013).** Housefly (*Musca domestica* L.) control potential of *Cymbopogon citratus* Stapf.(Poales: Poaceae) essential oil and monoterpenes (citral and 1, 8-cineole). *Parasitology research*, 112(1), 69-76.
- Lachman-White DA, Adams CD, Trotz UO (1992)** A guide to the medicinal plants of coastal Guyana. 2e éd Commonwealth Science council, Londre, p 92
- Leite, J., Maria De Lourdes, V. S., Maluf, E., Assolant, K., Suchecki, D., Tufik, S., ... & Carlini, E. A. (1986).** Pharmacology of lemongrass (*Cymbopogon citratus* Stapf). III. Assessment of eventual toxic, hypnotic and anxiolytic effects on humans. *Journal of ethnopharmacology*, 17(1), 75-83.
- Lima, E. O., Gompertz, O. F., Giesbrecht, A. M., & Paulo, M. Q. (1993).** In vitro antifungal activity of essential oils obtained from officinal plants against dermatophytes. *Mycoses*, 36(9-10), 333-336.
- Manosroi, J., Dhumtanom, P., & Manosroi, A. (2006).** Anti-proliferative activity of essential oil extracted from Thai medicinal plants on KB and P388 cell lines. *Cancer letters*, 235(1), 114-120.
- Mayaud L, Carricajo A, Zhiri A, et al. (2008)** Comparison of bacteriostatic and bactericidal activity of 13 essential oils against strains with varying sensitivity to antibiotics. *Lett Appl Microbiol* 47: 167–73
- Middleton B, Hui KP.** Inhibition of S-3-hydroxy-3-methylglutaryl-CoA reductase and *in vivo* rates of lipogenesis by amixture of pure cyclic monoterpenes [J]. *Biochem Pharmacol*, 1982, 31(18): 2897-2902.
- Ming, L. C., Figueiredo, R. O., Machado, S. R., & Andrade, R. M. C. (1995, August).** Yield of essential oil of and citral content in different parts of lemongrass leaves (*Cymbopogon citratus* (DC)

- STAPE) Poaceae. In *International Symposium on Medicinal and Aromatic Plants* 426 (pp. 555-559).
- Moreira, F. V., Bastos, J. F., Blank, A. F., Alves, P. B., & Santos, M. R. (2010).** Chemical composition and cardiovascular effects induced by the essential oil of *Cymbopogon citratus* DC. Stapf, Poaceae, in rats. *Revista Brasileira de Farmacognosia*, 20(6), 904-909.
- Nambiar VS, Matela H.** Potential functions of lemongrass (*Cymbopogon citratus*) in health and diseases [J]. *Int J PharmaceutBiol Arch*, 2012, 3(5): 1035-1043.
- Negrelle R.R.B. ; Gomes EC (2007)** *Cymbopogon citratus* (DC.) Stapf : Chemical Composition and Biological Activities. *Rev. Bras. Pl. Med.*, Botucatu, v.9, n.1, p.80-92
- Nogueira ACMA, Carvalho RR, Souza CAM, et al. (1995)** Study on the embryofeto-toxicity of citral in the rat. *Toxicology* 196: 105–13
- Ogunlana, E. O., Höglund, S., Onawunmi, G., & Sköld, O. (1986).** Effects of lemongrass oil on the morphological characteristics and peptidoglycan synthesis of *Escherichia coli* cells. *Microbios*, 50(202), 43-59. synthesis of *Escherichia coli* cells. *Microbios*, v. 50, n. 202, p. 43-59, 1987.
- Ohno T, Kita M, Yamaoka Y, et al. (2003)** Antimicrobial activity of essential oils against *Helicobacter pylori*. *Helicobacter* 8: 207–15
- Olivero-Verbel, J., Nerio, L. S., & Stashenko, E. E. (2010).** Bioactivity against *Tribolium castaneum* Herbst (Coleoptera: Tenebrionidae) of *Cymbopogon citratus* and *Eucalyptus citriodora* essential oils grown in Colombia. *Pest management science*, 66(6), 664-668.
- Olorunsanya AO, Olorunsanya EO, Bolu SAO, et al. (2010)** effect of graded levels of lemongrass (*Cymbopogon citratus*) on oxidative stability of raw or cooked pork patties. *Pakistan J Nutr* 9(5): 467–70
- Oloyede IO.** Chemical profile and antimicrobial activity of *Cymbopogon citratus* leaves[J]. *J Nat Prod*, 2009, 72: 98-103.
- Osmani, Z., & Sighamony, S. (1980).** Effects of certain essential oils on mortality and metamorphosis of *Aedes aegypti*. *Pesticides*, 14(9), 15-16.
- Pereira, R. P., Fachinetto, R., de Souza Prestes, A., Puntel, R. L., da Silva, G. N. S., Heinzmann, B. M., ... & Morsch, V. M. (2009).** Antioxidant effects of different extracts from *Melissa officinalis*, *Matricaria recutita* and *Cymbopogon citratus*. *Neurochemical research*, 34(5), 973-983.
- Phasomkusolsil, S., & Soonwera, M. (2011).** Efficacy of herbal essential oils as insecticide against *Aedes aegypti* (Linn.), *Culex quinquefasciatus* (Say) and *Anopheles dirus* (Peyton and Harrison). *Southeast Asian Journal of Tropical Medicine and Public Health*, 42(5), 1083.
- Puatanachokchai, R., Kishida, H., Denda, A., Murata, N., Konishi, Y., Vinitketkumnuen, U., & Nakae, D. (2002).** Inhibitory effects of lemon grass (*Cymbopogon citratus*, Stapf) extract on the early phase of hepatocarcinogenesis after initiation with diethylnitrosamine in male Fischer 344 rats. *Cancer letters*, 183(1), 9-15.
- Ritter RA, Monteiro MV, Monteiro FO, et al. (2012)** Ethnoveterinary knowledge and practices at Colaresisl and, Para state, eastern Amazon. *Brazil J Ethnopharmacol* 144: 346–52
- Sharma, P. R., Mondhe, D. M., Muthiah, S., Pal, H. C., Shahi, A. K., Saxena, A. K., & Qazi, G. N. (2009).** Anticancer activity of an essential oil from *Cymbopogon flexuosus*. *Chemico-biological interactions*, 179(2), 160-168.
- Shimoni, M. (1993).** Growth inhibition of plant pathogenic fungi by essential oils. *Hassadeh*, v.3, p.306-8 *Science & Emerging Technologies*, 8(2), 253-258.
- Singh P, Shukla R, Kumar A, et al. (2010)** Effect of citrus *reticulata* and *cymbopogon citratus* essential oils on *Aspergillus flavus* growth and aflatoxin production on *Asparagus racemosus*. *Mycopathologia* 170: 195–202
- Sosan, M. B., Adewoyin, F. B., & Adewunmi, C. O. (2001).** Larvicidal properties of three indigenous plant oils on the mosquito *Aedes aegypti*. *Nigerian Journal of Natural Products and Medicine*, 5(1), 30-33.
- Sousa SM, Silva PS, Viccini LF (2010)** Cytogenotoxicity of *Cymbopogon citratus* (DC) Stapf (lemon grass) aqueous extracts in vegetal test systems. *An Acad Bras Cienc* 82(2): 305–11
- Spring, M. A. (1989).** Ethnopharmacologic analysis of medicinal plants used by Laotian Hmong refugees in Minnesota. *Journal of ethnopharmacology*, 26(1), 65-91.

- Suman P.S. Khanuja\***, **Ajit K. Shasany**, **Anubha Pawar**, **R.K. Lal**, **M.P. Darokar**, **A.A. Naqvi**, **S. Rajkumar**, **V. Sundaresan**, **Nirupama Lal**, **Sushil Kumar (2004)**, Essential oil constituents and RAPD markers to establish species relationship in *Cymbopogon Spreng.* (Poaceae) *Biochemical Systematics and Ecology* 33 (2005) 171–186
- Tarkang PA**, **Agbor GA**, **Tsabang N, et al.** Effect of long-term oral administration of the aqueous and ethanol leaf extract of *Cymbopogon citratus* (DC. Ex Ness) Stapf.[J]. *Annals Biol Res*,2012, **3** (12): 5561-5570.
- Tchoumboungang F**, **Zollo PH**, **Dagne E, et al.** *In vivo* antimalarial activity of essential oils from *Cymbopogon citratus* and *Ocimum gratissimum* on mice infected with *Plasmodium Berghei* [J]. *Planta Med*, 2005, **71**(1): 20-23.
- Thangam, R.**, **Sathuvan, M.**, **Poongodi, A.**, **Suresh, V.**, **Pazhanichamy, K.**, **Sivasubramanian, S., ... & Kannan, S. (2014)**. Activation of intrinsic apoptotic signaling pathway in cancer cells by *Cymbopogon citratus* polysaccharide fractions. *Carbohydrate polymers*, *107*, 138-150.
- Trease, G.E.** A textbook of pharmacognosy. 9. ed. London: Bailière, Tindall and Cassell, 1996. p.201. TRIPPLEBROOKFARM. *Cymbopogon citratus*. Lemon grass. Available from:< <http://www.tripplebrookfarm.com/iplants/Cymbopogon.html> >. Accessed on: 20 jun. 2003. UNIVERSITY of Hawaii. Botany Department.
- Tzortzakis, N. G., & Economakis, C. D. (2007)**. Antifungal activity of lemongrass (*Cymbopogon citratus* L.) essential oil against key postharvest pathogens. *Innovative Food*
- Valarini, P. J., Frighetto, R. T. S., & Spadotto, C. A. (1996)**. Potential of the medicinal herbage *Cymbopogon citratus* for the control of pathogens and weeds in irrigated bean crop. *Científica (Jaboticabal)*, *24*(1), 199-214.
- Vigushin DM, Poon GK, Boddy A, et al. (1998)** Phase I and pharmacokinetic study of d-limonene in patients with advanced cancer. Cancer Research Campaign Phase I/II Clinical Trials Committee. *Cancer Chemother Pharmacol* 42: 111–7
- Vinitketkumnuen, U., Puatanachokchai, R., Kongtawelert, P., Lertprasertsuke, N., & Matsushima, T. (1994)**. Antimutagenicity of lemon grass (*Cymbopogon citratus* Stapf) to various known mutagens in salmonella mutation assay. *Mutation Research/Genetic Toxicology*, *341*(1), 71-75.
- Zheng, Guo Qiang, Kenney, Patrick M., et LAM, Luke KT.** Potential anticarcinogenic natural products isolated from lemongrass oil and galanga root oil. *Journal of Agricultural and Food Chemistry*, 1993, vol. 41, no 2, p. 153-156.
- Watanabe C, Hokari R, Komoto S, et al. (2010)** Lemon grass(*Cymbopogon citratus*) ameliorates murine spontaneous ileitisby decreasing lymphocyte recruitment to the inflamed intestine. *Microcirculation* 17: 321–32.